

KNOWLEDGE AND AWARENESS OF THE SYSTEMIC CONDITIONS AND ORAL HEALTH ASSOCIATION

Shahzeb Hasan Ansari

Riyadh Elm University, King Fahad Branch Road, Opposite passport office, Riyadh, KSA.

ABSTRACT

Introduction: Dental problems constitute the most among the majority of the world's population and its association with general medical health cannot be ignored. It is imperative for the medically compromised patients to take additional care of their oral cavity, which is generally ignored during the course of their treatment.

Materials and Methods: This is a cross sectional study, which involved the utilization of a closed ended questionnaire. Data collection was done as a combination of electronic. Paper survey Google forms were used among the general public to construct the electronic survey. All male and females above 18 years old were recruited. Collected data was analyzed using SPSS version 19.

Results: A total of N= 2234 Saudis took part in this study, which included 55% males and 45% females. As far as the age groups are concerned, 24% belonged to 18-30 years group, 32% to 31-45 years, 31% to 46-60 years and 13% to 60+ years. The mean scores of associations of cardiac diseases with oral health came out to be 1.18, for diabetes 2.13, for asthma 0.64, for liver disease 0.68 and for pregnancy 1.74.

Conclusion: Males showed a better level of knowledge as compared to females. Patients having a family member with or without a systemic condition showed varying responses; therefore, no distinctive association was discovered among these groups.

Key words: Systemic conditions, oral health attitudes, pregnancy related oral health.

Introduction

Health plays a critical role in determining the quality of any person's life, which includes general as well as oral health. Dental problems constitute the most among the majority of the world's population and its association with general medical health cannot be ignored¹⁻⁴. The medically compromised patients must take additional care of their oral cavity, which is generally ignored during their treatment. Chronic medical conditions including diabetes and obesity have been linked with poor oral health⁵⁻⁷.

Oral health serves as the most important determinant of health, which is of great concern among the middle as well as low socio-economic sectors. In support of the above-mentioned information regarding the quality of life, chronic oral diseases prominently affect this. Furthermore, oral health is seen to be poor among patients having chronic medical conditions^{8,9}.

Sohn & Rowe (2015) investigated the attitudes of medically compromised patients' parents, which revealed that the overall awareness was on the lower side¹⁰. These attitudes were comparatively better among the parents of healthy children. Pregnancy is also associated with compromised

oral health, which is evident in multiple studies. It is highly imperative to educate these groups of individuals as the levels of knowledge and attitudes below¹¹.

Patients with chronic liver disease are also known to be associated with having poor periodontal health. This is more commonly found in patients having a non-alcoholic fatty liver^{12, 13}. Similarly, patients with asthma have been associated with poor oral health due to the constant use of steroidal inhalers. Therefore, it is essential to educate such patients as their overall health-related behavior is unsatisfactory^{14, 15}.

Cardiovascular diseases have also been rumored to be linked with poor oral health, more specifically periodontal disease. However, Goulart et al (2017) revealed that this association was not proven when investigated a group of Brazilian patients¹⁶. However, patients with tuberculosis were found to have poor oral health, mainly due to their association with malnutrition¹⁷.

It is evident from the previous studies that even though the general public is aware of the association of systemic diseases with oral health, still the prevalence of poor oral health continues to be high. This has been observed when the cardiovascular disease was associated with poor oral health

and people's perception was inquired¹⁸. On the other hand, obesity and oral health have also been linked together and the knowledge among the people was found to be on the satisfactory side¹⁹. Whereas, knowledge of diabetes and oral health was also found to be satisfactory among the Pakistani population²⁰.

Aims of the study:

- To determine the knowledge and awareness of the general Saudi public towards the association of systemic diseases and oral health.
- To compare the knowledge based on gender, age, and educational level.

Materials and Methods:

This is a cross-sectional study, which involved the utilization of a closed-ended questionnaire. Data collection was done as a combination of electronic and paper surveys. Google forms were used to construct the electronic survey and distribute it among the general public. The questions included points related to the demographics, association of several systemic conditions with oral health including cardiac disease, diabetes, asthma, liver disease, and pregnancy. Data encoding was done giving the following values to responses when inquired about the above-mentioned systemic conditions:

- 0- Don't know
- 1- Not strong association
- 2- Somewhat strong association
- 3- Very strong association

Mean scores were being calculated for each association and mentioned in table 1. Convenient sampling was done. The sample size was decided to be a minimum of 1000, but we were able to achieve 2234 surveys filled by the participants.

Inclusions criteria:

All males and females above 18 years of age were recruited in this study.

Exclusion criteria:

Dental professionals.

Statistical analysis:

Collected data were subjected to analysis using SPSS version 19. Descriptive as well as inferential statistics were conducted using the Chi-square test with the value of significance kept under 0.05.

Results:

A total of N= 2234 surveys took part in this study, which included 55% males and 45% females (Figure 1). As far as the age groups are concerned, 24% belonged to the 18-30 years group, 32% to 31-45 years, 31% to 46-60 years, and 13% to 60+ years (Figure 2). We also divided the sample based on educational levels, which revealed that 8% were uneducated, 18% finished high school, 48% had a graduate degree and 26% had postgraduate qualification (Figure 3).

Further descriptive statistics showed that 64% of the participants reported having a family member having a systemic condition, 55% of them believed that there is an association of systemic conditions with oral health, 56% reported that the concerned physician is responsible to maintain the oral health of the patient. The mean scores of associations of cardiac diseases with oral health came out to be 1.18, for diabetes 2.13, for asthma 0.64, for liver disease 0.68, and pregnancy 1.74 (Table 1).

Table 2 shows the comparison of male and female participants on the responses they gave to survey questions. 66% of the females had at least one family member with a systemic condition as compared to 61% of males (p-value 0.024). 59% of males believed that there is an association of systemic conditions with oral health as compared to 49% of females (p-value: 0.021). 14% of male participants reported that cardiac diseases are strongly associated with oral health, whereas 18% of the females thought the same (p-value: 0.000). A strong association of diabetes with oral health was believed to exist according to 59% males and 51% females (p-value: 0.000). 46% of females thought that pregnancy is somewhat strongly associated with oral health as compared to 26% of males (p-value: 0.027).

We also compared our survey results based on age groups (Table 3), which reported that the younger age group (18-30 years) showed a higher inclination towards cardiac diseases (21%), diabetes (65%), and liver disease (21%) being strongly associated with effecting the oral health (p-values: 0.000, 0.005 and 0.000 respectively). The oldest age group (60+) believed asthma to be strongly associated with oral health as compared to other age groups (p-value: 0.002). As far as pregnancy and oral health is concerned, (31-45) years old study participants believed that there is a strong association between them (p-value 0.045).

Table 4 shows the comparison of survey responses according to the participants' family members having or not having systemic conditions (group A & B respectively). It was noted from the findings that 60% of group A participants believed that there is an association of systemic conditions with oral

health as compared to 48% of group B (p-value: 0.011). This table also shows varying responses to the association of different systemic conditions with oral health.




Figure 1: Gender ratio of the study participants




Figure 2: Age groups distribution of the study participants




Figure 3: Educational levels of the study participants

Survey Items	Descriptive Statistics
Family members having systemic conditions	Yes: 64% No: 36%
Association of systemic conditions with oral health?	Yes: 55% No: 18% Don't know: 26%
Association of cardiac diseases with oral health?	Mean score: 1.18
Association of diabetes with oral health?	Mean score: 2.13
Association of asthma with oral health?	Mean score: 0.64
Association of liver disease with oral health?	Mean score: 0.68
Association of pregnancy with oral health?	Mean score: 1.74
Is the person responsible to maintain oral health?	Dentist: 18% Concerned physician: 56% Patient himself: 26%

Table 1: Descriptive presentation of the responses from survey items

Mean score range: 0: Don't know – 3: Very strongly associated

Survey Items	Males	Females	P-value
Family members having systemic conditions	Yes: 61% No: 39%	Yes: 66% No: 34%	0.024
Association of systemic conditions with oral health?	Yes: 59% No: 15% Don't know: 26%	Yes: 49% No: 23% Don't know: 28%	0.021

Association of cardiac diseases with oral health?	Don't know: 40% Not strong: 34% Somewhat strong: 12% Very strong: 14%	Don't know: 25% Not strong: 20% Somewhat strong: 35% Very strong: 18%	0.000
Association of diabetes with oral health?	Don't know: 17% Not strong: 10% Somewhat strong: 8% Very strong: 59%	Don't know: 12% Not strong: 16% Somewhat strong: 21% Very strong: 51%	0.000
Association of asthma with oral health?	Don't know: 54% Not strong: 29% Somewhat strong: 7% Very strong: 9%	Don't know: 68% Not strong: 18% Somewhat strong: 10% Very strong: 5%	0.047
Association of liver disease with oral health?	Don't know: 62% Not strong: 19% Somewhat strong: 8% Very strong: 10%	Don't know: 64% Not strong: 16% Somewhat strong: 12% Very strong: 8%	0.134
Association of pregnancy with oral health?	Don't know: 24% Not strong: 18% Somewhat strong: 26% Very strong: 32%	Don't know: 10% Not strong: 15% Somewhat strong: 46% Very strong: 29%	0.027
Is the person responsible to maintain oral health?	Dentist: 15% Concerned physician: 59% Patient himself: 26%	Dentist: 21% Concerned physician: 50% Patient himself: 28%	0.000

Table 2: Comparison of survey responses based on gender

Survey Items	18-30 years	31-45 years	46-60 years	60+ years	P-value
Family members having systemic conditions	Yes: 76% No: 24%	Yes: 63% No: 37%	Yes: 42% No: 58%	Yes: 95% No: 5%	0.000
Association of systemic conditions with oral health?	Yes: 47% No: 32% Don't know: 21%	Yes: 61% No: 14% Don't know: 25%	Yes: 49% No: 14% Don't know: 37%	Yes: 71% No: 14% Don't know: 15%	0.000
Association of cardiac diseases with oral health?	Don't know: 30% Not strong: 35% Somewhat strong: 14% Very strong: 21%	Don't know: 44% Not strong: 29% Somewhat strong: 15% Very strong: 12%	Don't know: 49% Not strong: 23% Somewhat strong: 21% Very strong: 8%	Don't know: 38% Not strong: 15% Somewhat strong: 28% Very strong: 19%	0.000
Association of diabetes with oral health?	Don't know: 9% Not strong: 13% Somewhat strong: 14% Very strong: 65%	Don't know: 16% Not strong: 19% Somewhat strong: 8% Very strong: 57%	Don't know: 19% Not strong: 14% Somewhat strong: 12% Very strong: 54%	Don't know: 10% Not strong: 11% Somewhat strong: 35% Very strong: 44%	0.005
Association of asthma with oral health?	Don't know: 21% Not strong: 50% Somewhat strong: 13% Very strong: 16%	Don't know: 67% Not strong: 25% Somewhat strong: 2% Very strong: 6%	Don't know: 69% Not strong: 14% Somewhat strong: 12% Very strong: 15%	Don't know: 60% Not strong: 15% Somewhat strong: 8% Very strong: 17%	0.002

Association of liver disease with oral health?	Don't know: 32% Not strong: 34% Somewhat strong: 13% Very strong: 21%	Don't know: 77% Not strong: 10% Somewhat strong: 8% Very strong: 6%	Don't know: 69% Not strong: 14% Somewhat strong: 12% Very strong: 4%	Don't know: 41% Not strong: 10% Somewhat strong: 19% Very strong: 10%	0.000
Association of pregnancy with oral health?	Don't know: 21% Not strong: 32% Somewhat strong: 16% Very strong: 32%	Don't know: 14% Not strong: 17% Somewhat strong: 31% Very strong: 37%	Don't know: 10% Not strong: 12% Somewhat strong: 50% Very strong: 27%	Don't know: 34% Not strong: 17% Somewhat strong: 24% Very strong: 24%	0.045
Is the person responsible to maintain oral health?	Dentist: 34% Concerned physician: 55% Patient himself: 11%	Dentist: 24% Concerned physician: 49% Patient himself: 28%	Dentist: 19% Concerned physician: 54% Patient himself: 27%	Dentist: 18% Concerned physician: 29% Patient himself: 52%	0.000

Table 3: Survey response comparison based on age groups

Survey Items	Having a Condition (A)	Having No Condition (B)	P-value
Association of systemic conditions with oral health?	Yes: 60% No: 19% Don't know: 21%	Yes: 48% No: 17% Don't know: 37%	0.011
Association of cardiac diseases with oral health?	Don't know: 31% Not strong: 31% Somewhat strong: 19% Very strong: 20%	Don't know: 41% Not strong: 24% Somewhat strong: 26% Very strong: 9%	0.000
Association of diabetes with oral health?	Don't know: 10% Not strong: 19% Somewhat strong: 15% Very strong: 56%	Don't know: 23% Not strong: 7% Somewhat strong: 10% Very strong: 60%	0.000
Association of asthma with oral health?	Don't know: 61% Not strong: 25% Somewhat strong: 7% Very strong: 7%	Don't know: 56% Not strong: 24% Somewhat strong: 10% Very strong: 9%	0.006
Association of liver disease with oral health?	Don't know: 62% Not strong: 17% Somewhat strong: 11% Very strong: 12%	Don't know: 65% Not strong: 14% Somewhat strong: 16% Very strong: 5%	0.256
Association of pregnancy with oral health?	Don't know: 26% Not strong: 11% Somewhat strong: 31% Very strong: 33%	Don't know: 9% Not strong: 28% Somewhat strong: 35% Very strong: 28%	0.000
Is the person responsible to maintain oral health?	Dentist: 19% Concerned physician: 48% Patient himself: 33%	Dentist: 16% Concerned physician: 69% Patient himself: 16%	0.000

Table 4: Survey response comparison based on family members having or not having any systemic conditions

Discussion:

This study aimed to assess the Saudi public's knowledge regarding the association of systemic conditions with oral health. We were able to find significant comparisons among

the study groups based on their responses to the survey. We divided the sample into further subgroups according to their age, gender, educational levels, and having or not having a family member with a systemic condition. Diabetes mellitus is known to have a strong association with oral health, which

was elaborated by Jin et al (2016)²¹. According to them, 69% of the patients with diabetes showed poor oral health, which was reported by our study findings with the majority of the public being aware of this important association.

According to our study results (section 4), there is a healthy percentage of participants who are unaware of the strong association between systemic conditions and their effect on oral health. Similar findings were reported by Catteau et al (2016) when the study participants revealed a poor level of knowledge regarding the association of systemic conditions with oral health, especially in old, aged patients²².

Our study results have elaborated significant comparisons of the survey items based on gender, age, and participants having family members with or without systemic conditions (section 4). Their knowledge was compared, and it was found that all study groups exhibited a varying level of knowledge when it came to systemic health and its association with oral health. As far as gender was concerned, male participants slightly edged over the females when inquired about the above-mentioned association. However, these differences were statistically significant apart from the questions related to asthma and liver disease.

According to Mawardi, Elbadawi & Sonis (2015), it is the responsibility of both the dentist and the physician to maintain the oral health of their patients having systemic conditions²³. This is also found in our results as the majority of the participants voted in the favor of concerned physicians who should be careful when treating their patients having a certain systemic condition. Furthermore, the association of the cardiovascular disease with oral health cannot be ignored. Kjellstrom et al (2016) reported the above-mentioned association, which was not very well-acknowledged by the study participants of our investigation²⁴. It can be noted from our findings that the mean score was on the lower side when inquired about this association (section 4).

There is a scope of improvement in this study if we acquire information from the patients having systemic conditions. Their input would give this study a great deal of variation and uniqueness.

Conclusions:

In conclusion, overall knowledge of the study participants regarding the association of systemic conditions with oral health, males showed a better level of knowledge as compared to females, younger age group participants had a

better insight of this association as compared to middle-aged patients, patients having a family member with or without a systemic condition showed varying responses; therefore, no distinctive association was discovered among these groups.

References

1. Jahanmoghadam F, Gisour EF, Askari R, Rad M. Attitude Regarding Dental Stem Cells among Dental Practitioners in Kerman, Iran. *Arch. Pharm. Pract.* 2018;9(3):10-3.
2. Asgari, I., Soltani, S., Sadeghi, S. M. Effects of Iron Products on Decay, Tooth Microhardness, and Dental Discoloration: A Systematic Review. *Arch. Pharm. Pract.* 2020;11(1):60-82.
3. Kharalampos M, Put VA, Tarasenko SV, Reshetov IV. Comprehensive patient rehabilitation while performing immediate dental implant placement with the use of information-wave therapy (literature overview). *J. Adv. Pharm. Educ. Res.* 2020;10(4):1-6.
4. Bulgakova AI, Vasilyeva NA, Vasilyev EA. The clinical and immunological rationale for the use of prolonged action dental ointment in periodontology. *J. Adv. Pharm. Educ. Res.* 2019;9(4):65-69.
5. Sheiham, A. Oral health, general health, and quality of life. *Bulletin of the World Health Organization*, 2005; 83(9): 644.
6. Swapna, L., Koppolu, P., Prince, J. Oral health in diabetic and nondiabetic patients with chronic kidney disease. *Saudi Journal of Kidney Diseases and Transplantation*, 2017; 28(5): 1099.
7. Moore, P., Orchard, T., Guggenheimer, J., Weyant, R. Diabetes and oral health promotion: a survey of disease prevention behaviors. *The Journal of the American Dental Association*, 2000; 131(9): 1333-1341.
8. Petersen, P. The World Oral Health Report 2003: continuous improvement of oral health in the 21st century - the approach of the WHO Global Oral Health Programme. *Community Dentistry and Oral Epidemiology*, 2003; 31(s1): 3-24.
9. Munro, C.L., Grap, M.J. Oral health and care in the intensive care unit: state of the science. *American Journal of critical care*, 2004; 13(1): 25-34.
10. Sohn, H.A., Rowe, D.J. Oral health knowledge, attitudes, and behaviors of parents of children with diabetes compared to those of parents of children without diabetes. *American Dental Hygienists' Association*, 2015; 89(3): 170-179.
11. Rakchanok, N., Amporn, D., Yoshida, Y., Harun-Or-Rashid, M.D., Sakamoto, J. Dental caries and gingivitis among pregnant and non-pregnant women in Chiang

- Mai, Thailand. *Nagoya J Med Sci*, 2010; 72(1-2): 43-50.
12. Alazawi, W., Bernabe, E., Tai, D., Janicki, T., Kemos, P., Samsuddin, S., Syn, W.K., Gillam, D., Turner, W. Periodontitis is associated with significant hepatic fibrosis in patients with non-alcoholic fatty liver disease. *PloS one*, 2017; 12(12): p.e0185902.
 13. Akinkugbe, A.A., Slade, G.D., Barritt, A.S., Cole, S.R., Offenbacher, S., Petersmann, A., Kocher, T., Lerch, M.M., Mayerle, J., Völzke, H., Heiss, G. Periodontitis and Non-alcoholic Fatty Liver Disease, a population-based cohort investigation in the Study of Health in Pomerania. *Journal of clinical periodontology*, 2017; 44(11): 1077-1087.
 14. Bozejac, B.V., Stojšin, I., Đuric, M., Zvezdin, B., Brkanić, T., Budišin, E., Vukoje, K., Sečen, N. Impact of inhalation therapy on the incidence of carious lesions in patients with asthma and COPD. *Journal of Applied Oral Science*, 2017; 25(5): 506-514.
 15. Sharma, S., Gaur, P., Gupta, S., Kant, S. Impact of asthma on oral health: A review. *International Journal of Recent Scientific Research*. 2018; 9(5): 26512-26514.
 16. Goulart, A.C., Armani, F., Arap, A.M., Nejm, T., Andrade, J.B., Bufarah, H.B., Dezen, D.H.S. Relationship between periodontal disease and cardiovascular risk factors among young and middle-aged Brazilians. Cross-sectional study. *Sao Paulo Medical Journal*, 2017; 135(3): 226-233.
 17. Wu, L.L., Cheung, K.Y., Lam, P.Y.P., Gao, X. Oral health indicators for risk of malnutrition in elders. *The journal of nutrition, health & aging*, 2018; 22(2): 254-261.
 18. Sanchez, P., Everett, B., Salamonson, Y., Ajwani, S., Bhole, S., Bishop, J., Lintern, K., Nolan, S., Rajaratnam, R., Redfern, J., Sheehan, M. Oral health and cardiovascular care: Perceptions of people with cardiovascular disease. *PloS one*, 2017; 12(7), p.e0181189.
 19. Akram, Z., Abduljabbar, T., Hanif, A., Khan, A., Vohra, F. Attitude and knowledge of family medicine practitioners towards the association between periodontal disease and obesity. *Nigerian Journal of clinical practice*, 2017; 20(5): 595-599.
 20. Bangash, R.Y., Khan, A.U., Manzoor, M.A. Diabetic Patients; Level Of Awareness About Oral Health Knowledge, Attitude, And Practices. *Pakistan Oral & Dental Journal*, 2011; 31(2): 292.
 21. Jin, L.J., Lamster, I.B., Greenspan, J.S., Pitts, N.B., Scully, C., Warnakulasuriya, S. Global burden of oral diseases: emerging concepts, management, and interplay with systemic health. *Oral diseases*, 2016; 22(7): 609-619.
 22. Catteau, C., Piaton, S., Nicolas, E., Hennequin, M., Lassauzay, C. Assessment of the oral health knowledge of healthcare providers in geriatric nursing homes: additional training needs required. *Gerodontology*, 2016; 33(1): 11-19.
 23. Mawardi, H.H., Elbadawi, L.S., Sonis, S.T. Current understanding of the relationship between periodontal and systemic diseases. *Saudi medical journal*, 2015; 36(2): 150.
 24. Kjellstrom, B., Rydén, L., Klinge, B., Norhammar, A. Periodontal disease—important to consider in cardiovascular disease prevention. *Expert Review of Cardiovascular Therapy*, 2016; 14(9): 987–989.

Corresponding Author

Shahzeb Hasan Ansari

Riyadh Elm University, King Fahad Branch Road,
Opposite passport office, Riyadh, KSA.

Phone: +966534305435