CO-EXISTENCE OF ORAL CANCER AND OSMF IN DENTAL PATIENTS: A RETROSPECTIVE INSTITUTIONAL STUDY

Juhi Gupta¹, Anshul Aggarwal^{2*}, Rajendra Kumar Tewari³, Neha Agrawal⁴, Vipul Jain⁵

¹Assistant Professor, Department of Oral Pathology/Oral Medicine and Radiology, Dr. Z. A. Dental College, AMU Aligarh, India..
²Associate Professor & head, Department of Oral Pathology/Oral Medicine and Radiology, Dr. Z. A. Dental College, India..
³Professor, Department of Conservative Dentistry & Endodontics, Dr. Z. A. Dental College, AMU Aligarh, India.
⁴Associate Professor, Department of Periodontics and Community Dentistry, Dr. Z. A. Dental College, AMU Aligarh.
⁵Dental practitioner, India.

ABSTRACT

Introduction: Oral submucous fibrosis has been considered as a premalignant condition or potentially malignant disorder in the literature. It has been now considered an endemic disease in India & South Asian countries where gutka chewing, paan, or areca nut chewing has long been practiced. In recent years, concerns have increased in the new cases of OSMF with malignant transformation.

Aims and Objective: To evaluate the occurrence of oral cancer in OSMF patients and to understand the reason behind the increasing cases of oral cancer in patients suffering from OSMF.

Material and Method: 609 patients from the Western Uttar Pradesh population (01st March 2018 to 31st December 2019) who were reported to the Department of Oral Pathology/Oral medicine and radiology of our dental college with a clinical diagnosis of OSMF were included in the study. The male/female ratio was approximately 4.5:1. Patients with confirmed histopathological diagnosis of Oral cancer along with OSMF were considered as OSMF with oral cancer.

Result: Out of 609 OSMF patients who were screened for oral cancer, 196 cases were found positive for oral cancer. It means co-existence of oral cancer was found to be in 32.67% of cases, which was significant. All the variables like age, gender, socioeconomic status, and site of lesions were analyzed using the Chi-square test for proportion.

Conclusion: The co-existence of oral cancer in OSMF cases for a Western Uttar Pradesh Population was found to be significantly high. Easy accessibility to dried areca nut in attractive sachet or other forms of tobacco was one of the main causative factors. The need of the hour is to create awareness in the general public regarding the hazardous effect of Gutka chewing/tobacco/other forms especially in the young generation. The concerned authority must take essential steps to create awareness among the journal public about the same.

Key words: Oral Cancer, Areca nut, Gutka, OSMF, Arecoline.

Introduction

In 1966, Pindborg defined OSMF as "an insidious chronic disease that affects any part of the oral cavity and sometimes pharynx. Although sometimes associated with and/or preceded by vesicle formation, it is always associated with juxta-epithelial inflammatory response followed by fibroblastic changes in the lamina propria, with epithelial atrophy leading to the oral mucosa stiffness that causes trismus and difficulty in eating." ¹Initially, it was known as a precancerous condition but later on WHO has changed the nomenclature to a potentially malignant disorder.

Oral cancer is among the main causes of death globally. The prevalence of oral cancer is increasing day by day globally causing morbidity and mortality. The etiology is multifactorial and many etiological factors like genetic, host immunity, viral infection, or use of tobacco may lead to oral cancer 2,3 . Among all, the use of tobacco is the leading cause. In recent years there has been an exponential increase in the

number of oral cancer and the use of smokeless tobacco is one of the most important contributing factors ^{4,5}. Gutka that is made up of dried areca nut, slaked lime, tobacco, and some flavoring agents has become popular among the young generation and even females since few years. Easy accessibility and availability of this poison under the name of mouth freshener or breath freshener poise a great risk to health that is really alarming. Fresh mixture of Arecanut, Lime, Tobacco (Gutka) in different proportions, or as single/dual or triple mixture by an individual is also common in a certain section of the Indian population.

The use of areca nut alone or in combination with tobacco and flavoring agents not only increases the risk of oral cancer but also leads to the development of oral submucous fibrosis. Burning sensation for spicy food and inability to open the mouth interferes with the normal daily routine of the patient.

Many articles had been published in the past regarding the malignant potential of the OSMF. With this background, a

87

prospective study has been done on OSMF patients who visited the Department of Oral Pathology/Oral Medicine and Radiology, of our dental college in Aligarh; Uttar Pradesh; India.

Material and Methods

A cross-sectional survey was conducted among 609 patients who reported to the Department of Oral Pathology/Oral Medicine and Radiology of our Dental College from 01st March 2018 to 31st December 2019 with a clinical diagnosis of OSMF. The ethical clearance was obtained from the Institutional Research Committee; and informed consent was taken from each patient before starting the study. The male/female ratio was approximately 4.5:1.

The study tool was a questionnaire, which had the following parts:

- 1. Demographic details including Age, Gender, Education, Occupation, and Income
- 2. Questions related to use of tobacco type, frequency, and duration

Inclusion Criteria

- 1. Patients who had given consent to become a subject for the study
- 2. Patients with clinical and histopathological diagnosis of OSMF with or without oral cancer
- 3. Patients with a history of areca nut chewing, areca nut with tobacco, paan, paan masala, etc.

Exclusion Criteria

1. Patients who were not willing to give consent to become a subject for the study

- 2. Patients with oral cancer along with leukoplakic or erythroplakic patch
- 3. Patients with systemic conditions like chronic iron deficiency anemia
- 4. Oral ulcer with inconclusive histopathological diagnosis of oral cancer or lesion showing dysplasia.

Complete oral examination was done. Clinical diagnosis of OSMF was made based on palpable fibrous bands on buccal mucosa, restricted mouth opening/tongue movement, shrunken uvula, lip thinning, etc. Patients suffering from OSMF and also having non-healing ulcers were advised to undergo a histopathological examination of the ulcer. Patients with a confirmed histopathological diagnosis of oral cancer were given a diagnosis of OSMF with oral cancer. Most oral cancer patients were the cases referred from the Department of Radiotherapy JNMC for dental evaluation prior to radiation therapy for oral cancer.

The entire oral cavity was examined for oral ulcer, red or white patch; or lesions. Patients with confirmed histopathological diagnosis of oral cancer along with OSMF were considered as OSMF with oral cancer, the rest of the patients were kept on follow-up.

Results

Data obtained were tabulated and subjected to statistical analysis. The association of oral cancer was assessed with variables like age, gender, socioeconomic status, and site of a lesion using the Chi-square test for the proportion. For statistical analysis, SPSS v.22.0 was used.

	21 - 30	31 - 40	41 - 50	51 - 60	71 - 80	Total {Percentage (Actual
	years	years	years	years	years	Number)}
1 - 5 years	12 (3)	9.5 (7)	1.5 (1)	0	0	5.6 (11)
5 - 10 years	36 (9)	32.4 (24)	9.1 (6)	8.7 (2)	0	20.9 (41)
11 - 15 years	40 (10)	39.2 (29)	25.8 (17)	17.4 (4)	12.5 (1)	31.1 (61)
16 - 20 years	8 (2)	12.2 (9)	21.2 (14)	13.3 (3)	0	14.3 (28)
21 - 25 years	4 (1)	5.4 (4)	19.7 (13)	17.4 (4)	25 (2)	12.2 (24)
26 - 30 years	0	1.4 (1)	18.2 (12)	17.4 (4)	25 (2)	9.7 (19)
31 - 35 years	0	0	3 (2)	21.7 (5)	25 (2)	4.6 (9)
35 - 40 years	0	0	1.5 (1)	4.3 (1)	12.5 (1)	1.5 (3)
Total	100 (25)	100 (74)	100 (66)	100 (23)	100 (8)	100 (196)

Table 1. Age Distribution of OSMF and OSMF with oral cancer

The highest number of cases of OSMF with oral cancer was observed in the age group of 31-40 years. In addition, the highest cases of OSMF with oral cancer had the habit of consuming to bacco and other products for 11-15 years duration, which was statistically significant. (P >0.0001)



Figure 1. Different types of Smokeless Tobacco



Figure 2. Site Distribution of Oral Cancer



Figure 3: Socioeconomic Status and Concomitant presence of OSMF and Oral Cancer

Discussion

The burden of oral cancer is increasing in India every year. In the U.S. population, oral cancer accounts for approximately 3% of malignancies, however, it accounts for over 30% of all cancer (malignancy) cases in India ⁶. The site, pattern, and age distribution of oral cancer in India

differ considerably as compared to the Western population. Whereas in the Western population tongue is the common site of oral cancer, however In India it is more common in buccal mucosa or buccal vestibule. In our study also the most common site of occurrence of oral cancer with patient suffering with OSMF was found to be buccal mucosa figure 2 .

Oral submucous fibrosis is a known debilitating disease that interferes with everyday activity of a person as it causes restricted mouth opening and intolerance to spicy food. But now, the rising concern is the malignant potential of this condition. Many studies have been conducted by researchers in the past on different populations and they have reported different rates of malignant transformation of this disease in different populations and at different time frames for the same population. A study done by Pingborg et al. on 89 patients in Ernakulam District, Kerala showed oral cancer in 13% of patients with a malignant transformation rate of 4.5% over a period of 4 to 15 yr (median 8 yr)⁷. A study conducted by Murti et al on the same population for over 17 years showed a higher malignant transformation rate that is 7.6% as compared to 4.5%. ⁸

OSMF is characterized by metabolic alteration in collagen synthesis and destruction. The arecoline alkaloids present in areca nut leads to increased production of collagen and the collagen produced are resistant to destruction.

Gutka is a mixture of tobacco, spices, crushed areca nut (also called betel nut), and other ingredients. ⁹ Government has banned selling and purchase of gutka in India, however, consumption of commercially available areca nut with tobacco and other flavoring agents in different proportions has gained popularity among young Indian population.

Though it is a disease with a multifactorial origin, the development of this disease has shown dose-dependent relation with areca nut and betel quid chewing habit. In our study 91.1% study subjects had habit of chewing areca nut alone or in combination with tobacco. [Figure 1]

OSMF is a potentially malignant disorder and it increases the chance of development of oral cancer. Merchant et al. (2000) ¹⁰ in their study concluded that people with oral submucous fibrosis are 19.1 times more likely to develop oral cancer than those without it.

In our study, the incidence of oral cancer concomitant with oral submucous fibrosis was 32.67% which is higher (25.77%) than the incidence reported by Nishant Raj Chourasia et al. ¹¹ It may be attributed to the comparatively larger sample size of our study.

In our study, the maximum number of OSMF cases are seen in the age group of 21-30 yrs. [Table 1] The finding is consistent with the study done by Jha et al. 12

In our study, only 21% of patients were illiterate and 30% were either graduate or postgraduate. At least, 26% of patients had got education till high school. So, in the presence of formal education showed little effect on the general awareness regarding the health hazard of the areca nut chewing habit. But the low socioeconomic status is found to be related to higher incidence of OSMF with oral cancer. [Figure 3]. This finding of our study supports the findings of the study done by David I. Conway et al ¹³ that indicates that risk of oral cancer is significantly higher in patients belonging to low socioeconomic status.

Etiology- Reports regarding the contamination of areca nut with cancer-causing fungus (*Aspergillus flavus*), adulteration with several hazardous toxic metals, and even pesticides are freely available in the literature. Aflatoxins are fungal toxins produced by *A. parasiticus* and *Aspergillus flavus* have carcinogenic and several other adverse effects on human health.

The study done by Misra et al. & Muralimohan et al. (1981 & 1985)^{14, 15} found the fungus (*Aspergillus flavus* and *A. parasiticus*) in areca nut samples in the markets of Lucknow and Gulbarga cities of India. In a study done by Raisuddin et al. (1984)¹⁶, the presence of aflatoxin in areca nut in India was found to be in 37.5% of the sample with an average aflatoxin B1 content as high as 94 µg/kg and the average total aflatoxin (B1+B2+G1+G2) content of 137 µg/kg.

In India, the government regulation for the production and processing of areca nut and other flavoring agents under the name of mithi supari and other commercial products is not very stringent. As a result, the quality of the areca nut cannot be guaranteed.

Our study had a few limitations. As it was an institutional unicentric study, the outcomes cannot be generalized for the entire population. For more reliable outcomes, and generalization of the study findings, multicentre studies having a larger sample size need to be carried out. It was a cross-sectional study, so we can only assess the association of the presence of oral cancer in OSMF patients but cannot confirm that oral cancer developed secondary to OSMF.

Conclusion

OSMF is strongly associated with an increased risk of oral cancer, although the biology underlying this association has not yet been resolved. Chances of development of oral cancer in OSMF patients further increases due to the synergistic effect of other factors like genetic susceptibility, nutritional deficiency, and chronic trauma. The recent increased use of areca nut in combination with tobacco and other products has raised the number of new cases of oral cancer in patients suffering from OSMF and thus, the burden of oral cancer. The need of the hour is to increase the public awareness regarding the health hazard associated with chewing of areca nut or betel quid and stringent regulation by the government to the production of tobacco products.

References

- Ranganathan K, Mishra G. Review: An overview of classification schemes for oral submucous fibrosis. J Oral Maxillofac Pathol. 2006;10(2):55–8.
- Abualsaud NA, Alzahrani NS, Alsahli AA, Al Dahneem AH, Almohid RA, Alnattar AM, Alghorab LS, Nuwaylati NA, Aboulkhair RA, Abdulrhman M. Investigating the Effect of Cigarette Smoking on Gingival Diseases. Arch. Pharm. Pract. 2019;10(3):6-9.
- 3. Tadayon S, RaisiDehkordi Z, Jafarzadeh L. The Effect of Secondhand Smoke Exposure on Level of Maternal Thyroid Hormones. Int. j. pharm. phytopharm. res. 2018;8(5):53-8.
- Atik N, Umbarawan Y, Nurdiawan W, Avriyanti, E Achadiyani, Ruslami A. Aloe Vera protect the Rat's Lung after Cigarettes Smoke inducement: A histological study. J. Adv. Pharm. Educ. Res. 2019;9(3):96-100.
- AlShoubaki RE, AlZahrani AS. Outcomes of Gingival Depigmentation Among Smokers and Non-Smokers: A Comparative Study. Int. j. pharm. res. Allied sci. 2018;7(1):148-55.
- 6. Coelho KR. Challenges of the oral cancer burden in India. Journal of cancer epidemiology. 2012 Oct 4;2012.
- Pindborg JJ, Murti PR, Bhonsle RB, Gupta PC, Daftary DK, Mehta FS. Oral submucous fibrosis as a precancerous condition. Scand J Dent Res. 1984 Jun;92(3):224
- 8. Murti PR, Bhonsle RB, Pindborg JJ, Daftary DK, Gupta PC, Mehta FS. Malignant transformation rate in oral submucous fibrosis over a 17-year period. Community dentistry and oral epidemiology. 1985 Dec;13(6):340-1.
- 9. National Cancer Institute; https://www.cancer.gov/publications/dictionaries/canc er-terms/def/gutka
- 10. Merchant A, Husain SS, Hosain M, Fikree FF, Pitiphat W, Siddiqui AR, Hayder SJ, Haider SM, Ikram M, Chuang SK, Saeed SA. Paan without tobacco: an independent risk factor for oral cancer. International journal of cancer. 2000 Apr 1;86(1):128-31.
- 11. Chourasia NR, Borle RM, Vastani A. Concomitant association of oral submucous fibrosis and oral squamous cell carcinoma and incidence of malignant transformation of oral submucous fibrosis in a population of central India: A retrospective study. Journal of maxillofacial and oral surgery. 2015 Dec 1;14(4):902-6.
- 12. Jha RN, Kalyani PB, Savarkar SV. Incidence rate of oral submucous fibrosis (OSMF) and its etiology in patients visiting Government Dental College and Hospital, Jamnagar (GDCH, Jamnagar). Journal of Integrated Health Sciences. 2014 Jan 1;2(1):11.

- 13. Conway DI, Petticrew M, Marlborough H, Berthiller J, Hashibe M, Macpherson LM. Socioeconomic inequalities and oral cancer risk: a systematic review and meta-analysis of case-control studies. International journal of cancer. 2008 Jun 15;122(12):2811-9.
- 14. Misra A, Misra JK. Fungal flora of marketed betel nuts [Spoilage]. Mycologia. 1981.
- 15. MURALIMOHAN GU, REDDY CN. Incidence of Aflatoxins and Aspergillus flavus in marketed arecanut and soapnut powder in Gulbarga city. Indian Phytopathology. 1995;48(3):371-3.
- 16. Misra JK, Raisuddin S. Occurrence of fungi in marketed betel nut. Ind J Plant Path 1984;2:83-4.

Corresponding Author

Dr. Anshul Aggarwal

Associate Professor & head, Department of Oral Pathology/Oral Medicine and Radiology

Email Id: draggarwal.xray @ gmail.com

Phone Number: 7895093197